

INTERNATIONAL JOURNAL OF INSTITUTIONAL PHARMACY AND LIFE SCIENCES

Pharmaceutical Sciences

Review Article.....!!!

Received: 26-04-2016; Revised: 10-05-2016; Accepted: 11-05-2016

AN OVERVIEW ON NATURAL GUM AND ITS PHARMACEUTICAL APPLICATION

Darekar A.B.^{1*}, Desale K.B.¹, Saudagar R.B.²

¹Department of Pharmaceutics, R.G. Sapkal college of pharmacy, Anjaneri, Nashik-422213, Maharashtra, India.

²Department of Pharmaceutical chemistry, R.G. Sapkal college of pharmacy, Anjaneri, Nashik-422213, Maharashtra, India.

Keywords:

Natural Gum, biodegradable,
pharmaceutical excipients,
polysaccharides, NDDS

For Correspondence:

Dr. Darekar A.B.

Department of Pharmaceutics,
R.G. Sapkal college of
pharmacy, Anjaneri, Nashik-
422213, Maharashtra, India

E-mail:

kishor.desale9@gmail.com

ABSTRACT

The gums are widely used natural excipients for conventional and novel dosage forms. With the increasing interest in polymers of natural origin, the pharmaceutical world has compliance to use most of them in the formulations. In recent years, there has been a tremendous development in natural products, which are needed to be used for a variety of purposes. These natural materials have advantages over synthetic ones since they are chemically inert, nontoxic, less expensive, biodegradable and widely available. Gums are the potent candidates to be used in various pharmaceutical formulations as a potential candidate for novel drug delivery system (NDDS). In this review, we describe the developments in natural gums for use in the pharmaceutical sciences. They can also be modified in different ways to obtain tailor made materials for drug-delivery systems and thus can compete with the available synthetic excipients. Natural polymers are basically polysaccharides so they are biocompatible and without any side effects.

INTRODUCTION^[1,2]

The Excipients are additives used to convert the active pharmaceutical ingredients into dosage forms suitable for administration to patients.¹ Synthetic polymers offer a broad range of properties that can be reasonably well —built- in|| by design and modified by altering polymer characteristics. Plant products are therefore, attractive alternatives to synthetic products because of biocompatibility, low toxicity, environmental —friendliness and low price compared to synthetic products. Natural gums obtained from plants have diverse applications in drug delivery as a disintegrant, emulsifying agent, suspending agents and as binders. They have also been found useful in formulating immediate and sustained-release preparation.

Definition of Gums^[3]

The most common theories says that gums are formed as a natural phenomenon of the plant in which internal plant tissues disintegrate through a process called gummosis. This in turn form cavities, which exudes transformed carbohydrates called gums. Secondly it is caused as a result of injury to the bark or stem. Thirdly, some others attribute to fungi and bacteria attack to the plant. Majority of the gums are exuded from the stem. Only a few gums are obtained from roots, leaves and other parts of the plant. These gums on heating decompose completely without melting. Gums are found in large number of families. Notable among them are Leguminosae and Sterculiaceae. Other important gum yielding families are Anacardiaceae, Combretaceae, Meliaceae, Rosaceae and Rutaceae.

Classification of Gums^[4]

Gums are present in high quantities in a varieties of plants, animals, seaweeds, fungi and other microbial sources, where they perform a number of structural and metabolic functions; plant sources provide the largest amounts. The different available Gums can be classified as follows.

❖ **According to the charge**➤ **Anionic Polysaccharides**

- **Natural:** Alginic acid, pectin, Xanthan gum, Hyaluronic acid, Chondroitin sulfate, Gum Arabic, Gum Karaya, Gum Tragacanth
- **Semi-Natural:** Carboxymethyl, Chitin, Cellulose gum

➤ **Cationic Polysaccharides**

- **Natural:** Chitosan
- **Semi-Natural:** Cationic Guar gum.
- **Cationic-Hydroxyethylcellulose (HEC).**
- **NonionicPolysaccharides**
- **Natural:** Starch, Dextrins, Guar gum.
- **Semi-Natural:** Cellulose Ethers (e.g. hydroxyethyl cellulose, Methylcellulose, Nitrocellulose).
- **Amphoteric Polysaccharides**
- **Semi-Natural:**Carboxymethylchitosan, Nhydroxyl-Dicarboxyethylchitosan, Modified Potato starch.
- **Hydrophobic Polysaccharides**
- **Semi-Natural:**Cetylhydroxyethylcellulose, Polyquaternium.
- ❖ **According to the source**
- **Marine origin/algal (seaweed) gums:** Agar, Carrageenans, Alginic acid, Laminarin.
- **Plant origin**
- **shrubs/tree exudates**—Gum Arabica, GgumGhatti, Gum Karaya, Gum Tragacanth, Khaya and Albizia gums;
- **Seed gums**—Guar Gum, Locust bean Gum, Starch, Amylose, Cellulose
- **Extracts** -Pectin, Larch gum;
- **Tuber and roots**—Potato starch.
- **Animal origin:** Chitin and chitosan, Chondroitin sulfate, Hyaluronic acid.
- **Microbial origin (bacterial and fungal):** Xanthan, Dextrin, Curdian, Pullulan, Zanflo, emulsan, Baker's yeast glycan, schizophyllan, lentinan, krestin, scleroglucan.
- **Prepared gums**
- **Biosynthetic gums-** Xanthan, scleroglucan, dextrins.
- **Starch and its derivatives-**dextrins.
- **Cellulose derivatives.**
- **Semi-synthetic**
- **Starch derivatives** —Heta starch, Starch acetate, Sarch phosphates.

- **Cellulose derivatives** —Carboxymethyl cellulose (CMC), Hydroxyethyl cellulose, Hydroxypropyl methylcellulose (HPMC), methylcellulose (MC), Microcrystalline cellulose (MC).
- ❖ **According to shape**
 - **Linear:**Algins, Amylose, Cellulose, pectins.
 - **Branched**
 - **Short branches**—Xanthan, Xylan, Galactomannans;
 - **Branch-on-branch**—Amylopectin, Gum Arabic, Tragacanth.
- ❖ **According to Monomeric units in chemical structure**
 - **Homoglycans**— Amylose, Arabinanas, Cellulose;
 - **Diheteroglycans**— Algins, Carragennans, Galactomannans;
 - **Tri-heteroglycans**—Arabinoxylans, Gellan, Xanthan;
 - **Tetra-heteroglycans**—Gum Arabic, Psyllium seed gum;
 - **Penta-heteroglycans**—Ghatti gum, Tragacanth.

Advantages of Natural Gums in pharmaceutical science^[4]

- **Biodegradable**— Naturally available biodegradable polymers are produced by all living organisms. They represent truly renewable source and they have no adverse impact on humans or Environmental health (e.g. skin and eye irritation).
- **Biocompatible and non-toxic**—chemically, nearly all of these plant materials are carbohydrates composed of repeating sugar (monosaccharide's) units. Hence, they are non- toxic.
- **Low cost**—it is always cheaper to use natural sources. The production cost is also much lower compared with that for synthetic material. India and many developing countries are dependent on agriculture.
- **Environmental-friendly processing**—Gums from different sources are easily collected in different seasons in large quantities due to the simple production processes involved.
- **Local availability (especially in developing countries)** —in developing countries, government promote the production of plant like Guar gum and Tragacanth because of the wide applications in a variety of industries.

- Better patient tolerance as well as public acceptance-. There is less chance of side and adverse effects with natural materials compared with synthetic one. For example, PMMA, povidone.
- Edible sources—Most gums are obtained from edible sources.

Disadvantages of Natural Gums in pharmaceutical science^[4]

- Microbial contamination—the equilibrium moisture content present in the gums is normally 10% or more and, structurally, they are carbohydrates and, during production, they are exposed to the external environment and, so there is a chance of microbial contamination. However, this can be prevented by proper handling and the use of preservatives.
- Batch to batch variation—Synthetic manufacturing is a controlled procedure with fixed quantities of ingredients, while the production of gums is dependent on environmental and seasonal factors
- Uncontrolled rate of hydration—Due to differences in the collection of natural materials at different times, as well as differences in region, species, and climate conditions the percentage of chemical constituents present in a given material may vary. There is a need to develop suitable monographs on available gums.
- Reduced viscosity on storage—normally, when gums come into contact with water there is an increase in the viscosity of the formulations. Due to the complex nature of Gums (monosaccharide's to polysaccharides and their derivatives), it has been found that after storage there is reduced in viscosity.

Disadvantages of Synthetic Polymers in Pharmaceutical Sciences^[4,5,6,7,8]

- The synthetic polymers have certain disadvantages such as high cost, toxicity, environmental pollution during synthesis, non-renewable sources, side effects, and poor patient compliance.
- . Acute and chronic adverse effects (skin and eye irritation) have been observed in workers handling the related substances methyl methacrylate and poly- (methyl methacrylate).

- Reports of adverse reactions to povidone primarily concern the formation of subcutaneous granulomas at the injection site produced by povidone. There is also evidence that povidone may accumulate in organs following intramuscular injections.
- Acute oral toxicity studies in animals have indicated that carbomer-934P has a low oral toxicity at a dose of up to 8 g/Kg. Carbomer dust is irritating to the eyes, mucous membranes and respiratory tract. So gloves, eye protection and dust respirator are recommended during handling.
- Studies in rats have shown that 5% polyvinyl alcohol aqueous solution injected subcutaneously can cause anemia and can infiltrate various organs and tissues.
- Some disadvantages of biodegradable polymers used in tissue engineering applications are their poor biocompatibility, release of acidic degradation products, poor processing ability and rapid loss of mechanical properties during degradation. It has been shown that poly glycolides, polylactides and their co-polymers have an acceptable biocompatibility but exhibit systemic or local reactions due to acidic degradation products. An initial mild inflammatory response has been reported when using poly-(propylene fumarate) in rat implant studies.

Natural gums^[9,10]

Natural gums (gums obtained from plants) are hydrophilic carbohydrate polymers of high molecular weights, generally composed of monosaccharide units joined by glucosidic bonds. They are generally insoluble in oils or organic solvents such as hydrocarbons, ether, or alcohols. Gums are either water soluble or absorb water and swell up or disperse in cold water to give a viscous solution or jelly. On hydrolysis they yield arabinose, galactose, mannose and glucuronic acid.

Gums are produced by members of a large number of families but commercial exploitation is restricted to a few species of Leguminosae, and Combretaceae (Table 1). Gum is also extracted from seeds (Table 2), seaweeds (Table 3) microorganisms (Table 4), and *Aloe barbadensis* (aloe gum) wood chips of *Larix occidentalis* (stractan), seed coats or barns of corn, wheat, oats, barley, rice and soybean (Hemicellulose). Resins occur in a wide range of plants. They are formed in the specialized structures called ducts. With the exception of Lac produced by the Lac insect (*Laccifer lacca*) all the natural resins are of plant origin.

Table 1: List of a few plants, which are commercially tapped for, gums with their product names

Name of the source	Family	Exudate/ Product
<i>A. senegal</i> (L.) Willd.	Leguminosae	Gum Arabic
<i>Acacia seyal</i> Del.	Leguminosae	Gum Arabic
<i>Anogiessuslatifolia</i> Wall	Combretaceae	Gum ghatti
<i>Astragalusgummifer</i>	Leguminosae	Gum tragacanth
<i>Astragalusmicrocephalus</i> Willd.	Leguminosae	Gum tragacanth
<i>Azadirachtaindica</i> A. Juss.	Meliaceae	Neem gum
<i>Cochlospermumgossypium</i> L.	Cochlospermaceae	Gum karayaLannea

Table 2: List of plants which yield seed gum

Plant names	Family	Product
<i>Ceratonia siliqua</i> L. (Carob tree)	Leguminosae	Locust bean gum
<i>Cyamopsis tetragonolobus</i> (L.) Taub.	Leguminosae	Guar gum
<i>Plantago psyllium</i> L	Plantaginaceae	Pysillum seed
<i>Linum usitatissimum</i> L.	Linaceae	Flax seed
<i>Abelmoschus esculentus</i> (Pods)	Malvaceae	Okra gum
<i>Tamarindus indica</i>	Leguminosae	Tamarind gum

Table 3: List of a few seaweeds used as sources of gum

Plant names (Red Algae, Rhodophyceae)	Product
<i>Chondrus crispus</i>	Carrageenan
<i>C. ocellatus</i>	Carrageenan
<i>Gigartina stellata</i>	Carrageenan

<i>G. mamillosa</i>	Carrageenan
<i>G. acicularis</i>	Carrageenan
<i>G. radula</i>	Carrageenan
<i>G. pistillata</i>	Carrageenan
<i>Eucheumaspinosum</i>	Carrageenan or Agar
<i>E. muricatum</i>	Carrageenan or Agar
<i>E. cottonii</i>	Carrageenan or Agar
<i>E. edule</i>	Carrageenan or Agar
<i>Gelidiummamsii</i>	Agar
<i>G. cartilagineum</i>	Agar
<i>G. latifolium</i>	Agar
<i>P. densa</i>	Agar
<i>P. lucida</i>	Agar

Table 4: Biosynthetic gums (microbial gum)

Name of the organism	Product
<i>Xanthomonascompestris</i>	Xanthan
<i>Pseudomonas elodea</i>	Gellan
<i>Leuconostocmesenteroides</i>	Dextran
<i>Aureobasidiumpullulans</i>	Pullulan
<i>Hansenulahlolstii</i> (Yeast)	Phosphomannan - Y.2448
<i>Sclerotiumrolfsii</i> (Fungus)	Scleroglucan

Pharmaceutical Application of Natural Gum^[4,11]

Gums possess a complex, branched polymeric structure because of which they exhibit high cohesive and adhesive properties such properties used in pharmaceutical preparation. Hence gums find diverse application in pharmacy. They are ingredients in dental and other adhesive and as bulk laxative. These polymers are useful as tablets binder, disintegrating agent, emulsifier, suspending agent, thickener, gelling agent, stabilizing agent protective colloids in suspension and sustain agent in tablets. They act as adjuvant in some pharmaceutical formulation.

Table: 5 Pharmaceutical Application of natural gum

Common name	Botanical name	Family	Pharmaceutical name	Reference
Gum acacia	<i>Acacia Arabica</i>	<i>Leguminosae</i>	Binder, emollient, Suspending agent	50
Cashew gum	<i>Anacardium occidentale</i>	<i>Anacardiaceae</i>	Suspending agent	40-43
Xanthum gum	<i>Xanthomonas leucometris</i>	-	Suspending agent	72-74
Gum Tragacanth	<i>Astragalus Gummifer</i>	<i>Leguminosae</i>	Suspending, emulsifying agent	52
Albizi gum	<i>Albizia zizygia</i>	<i>Leguminosae</i>	Tablet Binder	34-36
sGuar gum	<i>Cyamopsis tetragonolobus</i>	<i>Leguminosae</i>	Binder, emulsifier, disintegrant	44-49
Khaya gum	<i>Khaya grandifolia</i>	<i>Meliaceae</i>	Binding agent	62

1. Application of gums in tablets formulation^[31,32,33,34,35]

Gums find its application in tablets formulation as a binder because of its adhesive nature. They impart cohesiveness to the powder mass and convert them into granules. They can also be used as disintegrates in tablets. the disintegrates property of gums due to absorb water and swell. They can swell up to 5 time their original volume this swelling lead to breakage of tablets into smaller particle which in turn improve dissolution rate. Example

- The binding agent *Butea monosperma* gum act as binder in ibuprofen tablets.
- The binding agent *Cassia roxburghii* seed as a binder in paracetamol tablets.
- The binding agent *Magnifera indica* gums as binder in paracetamol tablets.
- Cashew tree gums as binder in metronidazole tablets.
- Ziprasidon tablets based on dissolutions and dissolution efficiency value of performance of gums is as follows gum karaya> acacia>olibanum>tragacanth>guar gum.

2. Gums as emulsifying and suspending agent.^[36]

Gums act as emulsifying and suspending agent. They effectively stabilize the emulsion via interfacial absorption and subsequent formulation of condensed film of high tensile strength that resist coalescence of droplets. They stabilize oil/ water emulsion by forming strong multi molecular film round each oil globule thus retards the coalescence by hydrophilic barrier between oil and water phase.

Natural gums increase the hydration of the hydration layer around the suspended particle through hydrogen bonding and molecular interaction. Since this agent does not reduce the surface and interfacial tension, they function best in presence of the wetting agent. They also act as thickener and protective colloids. Natural gums are hydrophilic colloids which form dispersion with water and increase the viscosity of the continuous phase, so that solid particle suspended in it sufficient for long time to measure the uniform dose e.g. *Cordiagharaf* Gum.

3. Gums as sustaining materials in dosage form.^[37]

Gums can be use for sustaining the drug release. They have been used in tablets, suspension as or as matrix system for sustaining the drug release. This polymer when come and contact with water get hydrate and form a gel the drug release from this gel will be usually diffusion controlled hence release will be sustained over a long period e.g. guar gums, xanthenes gums and karaya gums.

4. Gums as coating agent.^[38]

Many gums act as coating agent, which can sustain the drug release, or can protect the drug from degradation in stomach as the number of coating increase the drug release is reduced e.g. *Grewia* Gum.

5. Application of gums in microencapsulation.^[39]

The gums because of their coating ability find application in microencapsulation of drug particles for sustaining the drug release. Gums from *Acacia nilotica* delile, *Acaicasenegal* wild and amizo gum has been studied for their microcapsulating properties using spray drying technique. Among these three *A. nilotica* is reported to be better microcapsulating agent e.g. gum kondagogu, gum Xanthan, gum guar.

6. Application of gums as gelling agent.^[40]

Gums can form a gel either alone or in combination with other. Gelling is a result of numerous inter and intra molecular association to produce three dimensional network, within which water

molecule are entrapped such association are brought about by Physical (pH change, altering temperature or chemical (addition of suitable reagent) treatment the mechanism of gelatin in acidic polysaccharides such as pectin is different. In this case the macromolecular chain is widely hydrogen bonded and as a result junction zone are formed between hydrogen bonded segments of chain. In Alginic acid, the gel formation occurs as result interaction with calcium ions. Galactomannan interact synergistically with xanthan gums and carrageenan to form as elastic gel e.g. locust bean gum.

Application of natural gum in novel drug delivery system

Table: 6 Application of natural gum in novel drug delivery system

Common name	Novel drug delivery system	Drug	Reference
Xanthan gum	Pellets Controlled drug delivery system	Diclofenac sodium. Theophylline	41,42
Sodium alginate	Bioadhesive microspheres	Gatifloxin. MetforminHCL	43,44
Karaya gums	Mucoadhesive and buccoadhesive	Nicotine	45
Guar gum	Colon targeted drug delivery	Albendazole Metronidazole methotrexate	46,47
Tamarind gum	Mucoadhesive drug delivery, Sustained releases	Diclofenac Sodium, Verapamil HCL	48,49
Acacia	Osmotic Drug Delivery	Water-insoluble naproxen	50,51
Gellan gums	Ophthalmic drug delivery, Beads, Floating in-situ gelling	Timolol, Propranolol, Amoxicillin	52,53,54
Locust bean gum	Controlled release agent	Nimodipine, Glipizide	55,56
Bhara gum	Microencapsulation	Famotidine	57
Cordia gum	Novel oral sustained release matrix forming agent in tablets. Suspension	Diclofenac Sodium, Paracetamol	58,59

CONCLUSION

Natural gum have advantages over synthetic ones since they are chemically inert, non-toxic, less expensive, biodegradable and widely available. The aim of this review article has been on the study of natural gum use in tablet formulation, advantages and disadvantages and various polymer used to design such dosage form. Natural gums are promising biodegradable polymeric materials. Many studies have been carried out in fields, including food technology and pharmaceuticals using gums. Various applications of gums have been established in the field of pharmaceuticals. However, there is a need to develop other natural sources as well as with modifying existing natural materials for the formulation of novel drug-delivery systems, biotechnological applications and other delivery systems. Therefore, in the years to come, there will be continued interest in natural gums and their modifications aimed at the development of better materials for drug delivery.

REFERENCES

1. Patel DM, Prajapat DG, Patel NM. Seed mucilage from *Ocimum americanum* as disintegrant in tablets: Separation and evaluation, Indian Journal of Pharmaceutical science. 2007; 69: 431-34.
2. The Joint IPEC – PQG Good Manufacturing Practices Guide for Pharmaceutical Excipients. 2006.
3. Market survey of gum and resin in India by RCDC (regional center for development co-operation center for forestry and governance. 2006:1-49.
4. Jani GK, Shah DP, Prajapati VD, V.C Jain *et al.* Gums and mucilage's: versatile excipients for pharmaceutical formulations. Asian Journal of Pharmaceutical Sciences. 2009; 4 (5): 308-322
5. Weller PJ, Owen SC, Polyvinyl alcohol. In: Raymond CR, Paul JS, Paul JW, Handbook of Pharmaceutical Excipients. 6th Ed, the Pharmaceutical Press and the American Pharmaceutical Association. 2003; 564-565.
6. Kottke KM, Edward MR. Tablet Dosage Forms. In: Banker GS, Rhodes CT. 3rd ed. Modern Pharmaceutics, New York, Marcel Dekker, Inc, 2002:287-333.
7. Aslam A, Parrott E, Effect of aging on some physical properties of hydrochlorthiazide tablets, Journal of Pharmaceutical sciences. 1971; 60: 263-266.
8. Avachat AM, Dash RR, Shrotriya S.N, Recent investigation of plant base natural gum, mucilage's and resins in novel drug delivery system, Indian journal of pharmaceutical education and research. 2011; 45(1):86-99.
9. Davison RL, Handbook of water-soluble gums and resins. New York, McGraw Hill Book Company; 1980.
10. Mantel CL, The water-soluble gums: Their botany source and utilization. Economic Botany. 1949; 3:3-31.
11. Giriraj Kulkarni T, Gowthamarajan K, Kumar M N, Suresh B, et al. therapeutics and pharmaceutical application. Natural Product Radiance. 2002.
12. Shefter E: Handbook of pharmaceutical excipients. Pharmaceutical Press, Edited by Raymond CR, Paul JS and Marian EQ, Edition 6, 2009: 703.
13. Ahmed BJ and Al-Ghazawi M: Sustained release characteristics of tablets prepared with mixed matrix of sodium carragennan and chitosan: Effect of polymer weight ratio, dissolution media and drug type. Drug Development and Industrial Pharmacy 2005; 31: 241-247.
14. Bonferoni MC, Rossi S, and Tamayo M: Employment of λ -carrageenan in a matrix system. Journal of Controlled Release 1994; 30: 175-182.

15. Pontes UR: Determination of HLB of Anacardium gum. Rev Farm Bioquim 1971; 2: 83-91.
16. Zakaria MB and Zainiah AR: Rheological properties of cashew gum. Carbohydrate Polymers 1996; 29: 2527.
17. Thierry N, George C and John F: Alginate and gellan gum as tablet coating. U.S. Patent No. 6326028 B1.
18. Kulkarni D, Dwivedi AK and Sarin JPS: Tamarind seed polyose: A potential polysaccharides for sustained release of verapamil hydrochloride as amodel drug. Indian Journal of Pharmaceutical Science 1997; 59: 1-7.
19. Dhopeswarkar V and Zatz JL: Evaluation of xanthan gum in the preparation of sustained release matrix tablets. Drug Development and Industrial Pharmacy 1993; 19: 999-1017.
20. Owen SC: Handbook of pharmaceutical excipients. Pharmaceutical Press, Edited by Raymond CR, Paul JS and Marian EQ, Edition 6, 2009: 782.
21. Ofoefule SI and Chukwu A: Application of Abelmoschus esculents gum as a mini-matrix for furosemide and diclofenac sodium tablets. Indian Journal of Pharmaceutical Science 2001; 63: 532-535.
22. John GL, Declan MD and James EK: The use of Agar as a novel filler for monolithic matrices produced using hot melt extrusion. European Journal of Pharmaceutics and Biopharmaceutics 2006; 64: 7581.
23. Oluwatoyin AO: Assessment of Albiziazygia gum as a binding agent in tablet formulations. ActaPharmaceutica 2005; 55: 263-276.
24. Pawar H and D'mello PM: Isolation of seed gum from *Cassia tora* and preliminary studies of its applications as a binder for tablets. Indian Drugs 2004; 41: 465-468.
25. Gowthamrajan K, Kulkarni GT and Muthukumar: Evaluation of Fenugreek mucilage as gelling agent. International Journal of Pharmaceutical Excipient 2002; 3: 16-19.
26. Kulkarni GT, Gowthamarajan K and Rao BG: Evaluation of binding property of *Plantagoovata* & *Trigonellafoenumgracecum* mucilage. Indian Drugs 2002; 39: 422-425.
27. Kale VV, Kasliwal R and Parida SK: Formulation and release characteristics of guar gum matrix tablet containing metformin HCl. International Journal of Pharmaceutical Excipient 2004; 2: 75-80.
28. Khullar P, Khar RK and Agrawal SR: Evaluation of guar gum in the preparation of sustained release matrix tablets. Drug Development and Industrial Pharmacy 1998; 24: 1095-1099.
29. Heda A and Shivhare U: Study of some natural hydrophilic polymers as matrix forming materials for sustained release tablet formulation. International Journal of Pharmaceutical Excipient 2004; 2: 69-74.
30. Odeku OA and Itiola OA: Evaluation of the effects of khaya gum on the mechanical and release properties of paracetamol tablets. Drug Development and Industrial Pharmacy 2003; 29: 311-320.
31. Deshmukh T, Patil P, Thakare V, Tekade B *et al*, evaluation of binding properties of *Buteamonosperma* Lam, Gum in tablet formulation, International journal of drug discovery and herbal research. 2011; 1(3): 128-133.
32. Arul Kumaran KSG, Palanisamy S, Rajasekaran A, *et al*. Evaluation of *Cassia roxburghii* seed gum as binder in tablet formulations of selected drugs. Int J Pharm Sci Nanotechnol. 2010; 2(4):726-732.
33. Singh AR, Selvam RP, Shivkumar T *et.al*, Isolation & characterization & formulation properties of natural Gum obtained from *Mangniferindica*, Int J Pharm Biomed Res. 2010; 1(2): 35-41.
34. Kwakye OK, Asantewaa Y and Kipo SM, Physiochemical and binding properties of Cashew Tree gum in Metronidazole tablet formulation, International Journal of Pharmacy and Pharmaceutical Sciences. 2010; 2(4): 105-109.
35. Prasanthi NL, Manikiran SS and Rao NR, Invitro drug release studies of Ziprasidone from tablets using natural gums from biosphere, Scholars Research Library. 2011; 3 (2):513-519.
36. Doharey V, Sharma N, Bindal MC, Assessment of the suspending properties of *Cordiagheraf* Gum on Paracetamol suspension, Scholars Research Library. 2010; 2 (1): 510-517.
37. Jian S, Yadav SK, Patil UK, Preparation and Evaluation of Sustained Release Matrix Tablet of Furosemide, Research journal of pharmacy and technology. 2008; 1(4): 374-376.
38. Ogaji IJ, Okafor IS, and Hoag SW, Grewia Gum as a Film Coating Agent in Theophylline Tablet Formulation: Properties of Theophylline tablets coated with grewia gum as a film coating agent, Journal of Pharmacy and Bioallied Science. 2013; 5(1): 53-60.

- 39.Mankala SK, Nagamalli NK, Raprla R, Kommula R, Preparation and Characterization of Mucoadhesive Microcapsules of Gliclazide with Natural Gums, Stamford journal of Pharmaceutical sciences. 2011; 4(1): 38-48.
- 40.Dionísio M and Grenha A, Locust bean gum: Exploring its potential in biopharmaceutical application, Journal of pharmacy and Bioallied sciences. 2012; 4(3): 175-185.
- 41.Santos H, Veiga F, Pina M.E, Sousa J.J. Compaction compression and drug release properties of diclofenac sodium and ibuprofen pellets comprising xanthan gum as a sustained release agent. International journal of Pharmaceutics, 2005; 295(1-2):15-27.
- 42.Vendruscolo CW, Andreazza IF, Ganter JL, *et al.* Xanthan and galactomannan (from *M. scabrella*) matrix tablets for oral controlled delivery of theophylline, International journal of Pharmaceutics, 2005; 296(1-2):1-11.
- 43.Kesavan K, NathG, Pandit J.K, sodium alginate based mucoadhesive system for gatifloxacin and its in vitro antibacterial activity, scientiapharmaceutica. 2010; 78(4):341-357.
- 44.Balasubramaniam J, Rao VU, Vasudha M, Babu J, Rajinikanth PS, Sodium alginate microspheres of metformin HCl: formulation and in vitro evaluation, current drug delivery, 2007; 4(3):249-56.
- 45.Park C.R, Munday DL. Evaluation of selected polysaccharide excipients in buccoadhesive tablets for sustained release of nicotine, Drug Development and industrial pharmacy. 2004; 30(6): 609-617.
- 46.Krishnaiah YSR, *et al.* Development of colon targeted oral Guar gum matrix tablets of Albendazole for the treatment of helminthiasis. Indian journal of pharmaceutical science.2003; 65(4):378-385.
- 47.Krishnaiah YSR *et al.* Guar gum as a carrier for colon specific delivery; influence of Metronidazole and Tinidazole on In Vitro release of Albendazole from Guar gum matrix tablets. Journal of Pharmacy and Pharmaceutical Science, 2001; 4(3):235-243.
- 48.Kulkarni G.T, Gowthamarajan K, Dhobe R.R, *et al.*, Development of controlled release spheroids using natural polysaccharide as release modifier, Drug Delivery. 2005;12(4): 201-206.
- 49.Kulkarni D, Dwivedi A.K, Sarin J.P.S, Singh S, Tamarind Seed Polyose : A Potential Polysaccharide For Sustained Release Of Verapamil Hydrochloride As a Model Drug, Indian journal of pharmaceutical sciences. 1997; 59(1):1-7.
50. Lu EX, Jiang, Zhang QZ, *et.al.* A water-insoluble drug monolithic osmotic tablet system utilizing gum Arabic as an osmotic, suspending and expanding agent. Journal of control release Controlled Release. 2003; 92(3): 375-382.
- 51.Beneke CE, Viljoen AM, Hamman JH, Polymeric plantderived excipients in drug delivery Molecules. 2009; 14: 26022620.
- 52.Rozier A, Mazuel C, Grove J, Plazonnet B, and Functionality testing of gellan gum: a polymeric excipient material for ophthalmic dosage forms. International journal of Pharmaceutics. 1997; 153(2):191-198.
- 53.Kedzierewicz F, Lombry C, Rios R, *et al.* Effect of the formulation on the in-vitro release of propranolol from gellan beads, International journal of pharmaceutics, 1999;178(1):129136.
- 54.Rajinikanth PS, Balasubramaniam J, Mishra B, Development and evaluation of a novel floating in situ gelling system of amoxicillin for eradication of *Helicobacter pylori*, International of pharmaceutics, 2007;335(1):114-122.
- 55.Bangale GS , *et.al.* Formulation and Evaluation of Natural Gum Based Matrix Tablets for Oral Controlled Delivery of Nimodipine, Indian Journal of Pharmaceutical Education and Research, 2011;45(4):375-383.
- 56.Xiaohong MG, Michae JT, John NS, Influence of physiological variables on the in-vitro drug-release behavior of a polysaccharide matrix controlled-release system, Drug Development and industrial pharmacy. 2003; 29(1):19-29.
- 57.Nayak BS, Nayak UK, Patro KB, *et al.* Design and evaluation of controlled release Bhara gum microcapsules of famotidine for oral use, Research Journal of Pharmacy and technology. 2008; 1(4):433-437.
58. Mukherjee B, Dinda SC, Barik BB, Gum Cordia: A novel matrix forming material for enteric resistant and sustained drug delivery - A Technical Note, AAPS Pharm SciTech, 2008;9(1):330-333.
- 59.Doharey V, Sharma N, Bindal MC, Assessment of the suspending agent of Cordial gheraf gum on paracetamol suspension. Scholars research library.2010; 2(1):510-517.